

Particle Acceleration, Magnetic Field Generation, and Emission in Relativistic Shocks

Ken Nishikawa (NSSTC/MSFC) (homepage: <http://gammaray.nsstc.nasa.gov/~nishikawa>), C. Hededal (Niels Bohr Inst.)

P. Hardee (Univ. of Alabama, Tuscaloosa), G. Richardson (UAH), R. Preece (NSSTC/MSFC), H. Sol (OPM), G.J. Fishman (NSSTC/MSFC)

Abstract

Shock accelerations are ubiquitous phenomena in astrophysical plasmas. Plasma waves (e.g., Buneman instability, two-streaming, Weibel instabilities) created in the shock are responsible for particle (electron, positron, and ion) acceleration. Using a 3-D relativistic electromagnetic particle (REMP) code, we have investigated particle acceleration in relativistic jets¹. Simulations show that the **Weibel instability** created at the shock front accelerates particles perpendicular and parallel to the jet propagation direction. This instability is also responsible in generating magnetic fields in the relativistic jets. The simulation results show that the growth rate of the Weibel instability depends on the size of the jet, composition of the jet, and the orientation and strength of ambient magnetic fields. The magnetic fields generated by the Weibel instability create highly nonuniform, small-scale magnetic fields, which contribute to the electron's transverse deflection. The radiation by deflected electrons is different from synchrotron radiation, which is called **jitter radiation**.

Scientific objects

- How do shocks in relativistic jets evolve in **accelerating particles and emission**?
- Do **reverse shocks** create **internal shocks** ?
- How do **3-D relativistic particle simulations** reveal the dynamics of shock front and transition region?
- What is the **main acceleration mechanism** in relativistic jets, **shock surfing, wakefield, Fermi models or stochastic processes**?
- How do shocks in **relativistic jets** evolve in the **different ambient plasma and magnetic field conditions** in various astrophysical phenomena?
- How does **Weibel instability** contribute to jitter radiation?


¹ Nishikawa et al. 2003, ApJ, 595, 555, Particle acceleration in relativistic jets due to Weibel instability (astro-ph/0305091)

Necessity of particle simulation for particle acceleration

- MHD simulations provide **global dynamics** of relativistic jets including hot spots
- MHD simulations include **heating** due to shocks, however do not create high energy particles
- In order to take account of acceleration the **kinetic effects** need to be included
- **Test particle (Monte Carlo) simulations** can include kinetic effects, but not self-consistently
- **Particle simulations provide particle acceleration and emission self-consistently**, however due to the computational limitations, the size of jet is small comparing with MHD simulations

Schematic GRB from a massive stellar progenitor


(Meszaros, Science 2001)



Collisionless shock


Electric and magnetic fields created self-consistently by particle dynamics randomize particles

(Buneman 1993)



$$\begin{aligned} \partial B / \partial t &= -\nabla \times E \\ \partial E / \partial t &= \nabla \times B - J \\ dm_{\alpha} v / dt &= q(E + v \times B) \\ \partial \Delta / \partial t + \nabla \cdot J &= 0 \end{aligned}$$

Weibel instability



Flat jet injected to unmagnetized plasma

Electron-positron jet, $m_p/m_e = 1$

$\beta = v_j/c = 0.9798$, $v_{ej}/c = 0.1$

$\eta = n_j/n_a \approx 0.741$

$\gamma = (1 - (v_j/c)^2)^{-1/2} = 5$


$v_{je} = 3v_{ej}$, $v_{jp} = 3v_{pj}$

$\omega_{pe} \Delta t = 0.013$

$\lambda_{ce} = c/\omega_{pe} = 9.6 \Delta$, $\lambda_e = v_{ej}/\omega_{pe} = 0.96 \Delta$


Electron density (arrows: B_z, B_x)

$\omega_{pe} t = 23.4$



Perpendicular current J_y ($J_{z,x}$)


$\omega_{pe} t = 23.4$



Evolution of B_x due to Weibel instability


(convective instability)

$\omega_{pe} t = 23.4$




Comparison between ele-ion and ele-pos

$\omega_{pe} t = 23.4$



Density and J_z in x-y plane

$\omega_{pe} t = 23.4$



Summary

- Simulation results show **Weibel instability** which creates filamented currents and density along the propagation of jets.
- The growth rate of Weibel instability depends on the **size of jets, compositions, strength and direction of ambient magnetic fields**.
- In a one-dimensional system **Buneman instability** is responsive to the surfing acceleration, **Weibel instability** is excited in the 3-D system.
- In order to understand the complex shock dynamics of relativistic jets, further simulations with additional physical mechanisms such as **radiation loss and inverse Compton scattering** are necessary.
- The magnetic fields created by Weibel instability generate highly inhomogeneous magnetic fields, which is responsible for **Jitter radiation** (Medvedev, 2000, ApJ).
- Weibel instability may play a major role in particle acceleration in relativistic jets
- **The dynamics of jet head is complicated** and further investigation is necessary

HEAD Meeting, New Orleans, September 8 – 11, 2004

Future plans for particle acceleration in relativistic jets

- **Further simulations** with a systematic parameter survey will be performed in order to understand shock dynamics
- In order to investigate shock dynamics **further diagnostics** will be developed
- In order to improve the performance of the code, **HPF** or **MPI** will be used
- Implement **better boundary conditions** at the free boundaries
- Investigate **synchrotron (jitter) emission** from the accelerated electrons and compare with observations (Blazars and gamma-ray burst emissions)
- Develop a new code implementing **synchrotron loss and/or inverse Compton scattering**
- Compare simulation results with **relativistic electron-positron experiments at SLAC** to understand particle acceleration in astrophysical relativistic jets

Electron velocity distributions and phase space

